version 9.11.2011

arXiv:1104.2574v3 [hep-ph] 9 Nov 2011

Generalisation of QED including scalar boson coupling

H.P. Morsch[1]
National Centre for Nuclear Research, Pl-00681 Warsaw, Poland

Abstract

Quantum electrodynamics with massless fermions (quantons) has been extended to include scalar coupling between two boson fields. New features are observed: 1. A term with derivative coupling appears in the Lagrangian, which leads to the long sought confinement potential known in hadronic systems. 2. Divergences in boson-exchange matrix elements disappear, giving rise to solutions in full space. 3. The scalar boson coupling allows a reduction of the formalism to three dimensions, from which the mass and static properties of stationary systems can be extracted. Application of this model to the hydrogen atom and positronium as well as to the lowest scalar meson $\sigma(600)$ yields a quantitative description of basic properties of these systems.

PACS/ keywords: 11.15.-q, 12.40.-y, 14.40.Cs, 31.15.Ne/ Extension of QED by scalar coupling between gauge bosons. Confinement and boson-exchange potential. Application to the hydrogen atom and positronium, structure of the $\sigma(600)$ meson.

In spite of large successes of the Standard Model of particle physics [1] (SM), constructed from different gauge theories, quantum electrodynamics, weak interaction theory, and quantum chromodynamics, this model is not really satisfactory in the understanding of nature. There are severe problems with gravitation, which cannot be included in the SM, (Dirac) neutrino masses are zero in conflict with neutrino oscillation results, and further the flavour families of leptons and hadrons are not understood. Therefore, extensions of the SM are of large current interest. Among these models grand unified theories, in particular of supersymmetric structure (SUSY) have received special attention, see e.g. the discussion in ref. [1]. In SUSY new particles are predicted with large masses,

---

[1] postal address: Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich, Germany
E-mail: h.p.morsch@gmx.de

which are currently subject of intense searches in high energy experiments at Tevatron and LHC ref [2]. But so far evidence for these particles has never been found.

In the present paper a very different generalisation of a Lagrangian for fundamental forces is presented, the inclusion of scalar coupling of elementary boson fields in quantum electrodynamics (QED) with massless[2] fermions (quantons, q). This approach is guided by experimental observations of strong scalar fields in hadron structure [3, 4] and scattering [5, 6], in particular at high energies, where the cross section is dominated by (scalar) Pomeron-exchange [7]. Also, such a Lagrangian allows a description of static properties of stationary states, as radii and form factors, in a fully relativistic framework.

To take into account the scalar boson degree of freedom in QED, the Lagrangian has to include in addition to the gauge boson field A^μ a scalar coupling of two boson fields $A_\mu A^\mu$, which leads to $\mathcal{L} = \frac{1}{\tilde{m}^2}\bar{\Psi}\ i\gamma_\mu D^\mu (D_\nu D^\nu)\Psi\ -\ \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$, where $\tilde{m}$ is a mass parameter and Ψ a massless fermion (quanton, q) field. This is generally a two component field with charged and neutral quantons, but here only charged quantons are considered with charge coupling $g = g_e$. The covariant derivative is given by $D_\mu = \partial_\mu - igA_\mu$ and the field strength tensor by $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$.

This Lagrangian includes higher derivatives of the fermion fields through gauge invariant scalar boson coupling. Such Lagrangians are known to lead to severe problems, if they are not constrained, see eg. ref. [8]. If fields and their derivatives appear in the Lagrangian, Heisenberg's uncertainty principle may not be respected, resulting in unphysical solutions, which may be unbound from below, violate unitarity and give rise to ghosts. Unphysical solutions can be avoided, if by appropriate physical constraints only fields or their derivative appear, but not both.

In the above Lagrangian one may introduce a symmetry between fermions and one boson field

$$D^\nu = (\partial^\nu - igA^\nu) = constant = 1\ , \tag{1}$$

which yields

$$\partial(\partial^\nu - igA^\nu) = \partial\partial^\nu - ig\partial A^\nu = 0. \tag{2}$$

Eq. (1) gives rise to a constrained Lagrangian in the form

$$\mathcal{L} = \frac{1}{\tilde{m}}\bar{\Psi}\ i\gamma_\mu D^\mu D_\nu \Psi\ -\ \frac{1}{4}F_{\mu\nu}F^{\mu\nu}\ . \tag{3}$$

---

[2] with a negligible mass with respect to the mass scale of particle physics

The first term of the Lagrangian (3) can be written by

$$\mathcal{L}_1 = \frac{1}{\tilde{m}}\bar{\Psi}\ i\gamma_\mu(\partial^\mu - igA^\mu)(\partial_\nu - igA_\nu)\Psi$$

$$= \frac{1}{\tilde{m}}\bar{\Psi}\ i\gamma_\mu[\partial^\mu\partial_\nu - igA^\mu\partial_\nu - igA^\mu\partial_\nu - g^2A^\mu A_\nu]\ \Psi\ . \tag{4}$$

Eq. (2) shows that the two first terms in the bracket cancel. By inserting the relation (1) in inverse form one obtains

$$\mathcal{L}_1 = \frac{1}{\tilde{m}^2}\bar{\Psi}\ i\gamma_\mu(\partial^\nu - igA^\nu)[-igA^\mu\partial_\nu - g^2A^\mu A_\nu]\ \Psi$$

$$= \frac{1}{\tilde{m}^2}\bar{\Psi}\ i\gamma_\mu[-ig(\partial^\nu\partial_\nu - ig\partial^\nu A_\nu)A^\mu - g^2A^\mu\partial_\nu A^\nu + ig^3A^\mu A_\nu A^\nu]\ \Psi\ . \tag{5}$$

Again the first two terms cancel and the Lagrangian (5) reduces to the following two terms

$$\mathcal{L}_{2g} = \frac{-ig^2}{\tilde{m}^2}\ \bar{\Psi}\gamma_\mu A^\mu(\partial_\nu A^\nu)\Psi \tag{6}$$

and

$$\mathcal{L}_{3g} = \frac{-g^3}{\tilde{m}^2}\ \bar{\Psi}\gamma_\mu A^\mu(A_\nu A^\nu)\Psi\ . \tag{7}$$

These two pieces of the Lagrangian $\mathcal{L}_1$ do not have derivatives of the fermion field. Further, the derivative of A^ν appears in $\mathcal{L}_{2g}$, but this field appears in $\mathcal{L}_{3g}$ only in combination with another boson field. Therefore, the problems of higher derivative Lagrangians are avoided. This is confirmed in the application to atomic and hadronic bound states, which leads to stable physical solutions, as discussed below, without ghost solutions or violation of unitarity. Concerning the renormalisability of the Lagrangian, which is generally not possible, if a mass factor with exponent ≥ 1 appears in the denominator, it will be shown that the present Lagrangian leads to finite matrix elements and therefore renormalisation is not needed.

To study bound state properties arising from the above Lagrangians (6) and (7), ground state matrix elements of field operators K have to be calculated. First order matrix elements may be written in the form $< g.s.|\ K(\tilde{p}'-\tilde{p})\ |g.s. >\sim \bar{\psi}(\tilde{p}')\ K(\tilde{p}'-\tilde{p})\ \psi(\tilde{p})$, where $\psi(\tilde{p})$ is a two-quanton wave function $\psi(\tilde{p}) = \frac{1}{\tilde{m}^3}\Psi(p)\Psi(k)$ and $K(\Delta\tilde{p})$ a kernel directly related to the multi-boson field operators of the Lagrangians (6) and (7). In principle, a Lagrangian with three-boson coupling could lead to a complicated structure of the kernel with propagation of three bosons. However, the fact that the field symmetry (1) requires a velocity distribution of the quanton field within the product of two boson field, the two-boson field can be regarded as a background field $w(q)$, in which for the case of $\mathcal{L}_{3g}$ a vector boson propagates. This is also important for the derivation of a geometric boundary condition (see below). Further, this avoids problems with causality and allows to go back to QED for all problems not directly connected to bound states.

In the language of Feynman diagrams the background field $w(q)$ may be viewed as two boson lines emitted and reabsorbed in the quanton field, leading only to a renormalisation of the wave function. $\mathcal{L}_{2g}$ includes only two boson fields and is interpreted in the Feynman picture as a $q\bar{q}$ contact term.

Interestingly, the Lagrangian $\mathcal{L}_{2g}$ does not appear in normal gauge theories. The corresponding kernel is dynamically generated by the two-boson field $w(q)$. By applying the derivative operator $\partial w(q)$ twice, a matrix element is obtained in the form

$$\mathcal{M}_{2g} = \frac{\alpha^2}{\tilde{m}^5} \bar{\psi}(\tilde{p}')\gamma_\mu \ \partial^\mu w(q) \partial^\rho w(q) \ g_{\mu\rho} \ \gamma_\rho \psi(\tilde{p}) \tag{8}$$

with $\alpha = g^2/4\pi$ and $\tilde{p}' - \tilde{p} = q$. By contraction of the γ-matrices to $g_{\mu\rho} = \frac{1}{2}(\gamma_\mu\gamma_\rho + \gamma_\rho\gamma_\mu)$ [9] and reduction to three dimensions [10], $\mathcal{M}_{2g}$ may be written by

$$\mathcal{M}_{2g} = \bar{\psi}(\tilde{p}') \ T_{2g}(q) \ \psi(\tilde{p}) \tag{9}$$

with $T_{2g}(q) = \frac{\alpha^2}{\tilde{m}^3} \ (dw(q)/dq)^2$, which appears to be a kinetic energy term. Transformed to coordinate space x, this term can be identified with the kinetic energy of a Hamiltonian with the form

$$-\frac{\alpha^2 \hbar^2 \tilde{E}^2}{2\tilde{m}^3} \ \nabla^2 w(x) + V_{2g}(x) \ w(x) = E_i \ w(x) \ , \tag{10}$$

where w(x) is the two-boson wave function and $\tilde{E}$ the mean energy of the scalar field. From this Hamiltonian, the potential $V_{2g}(x)$ can be calculated by assuming that the lowest eigenstate E_o is the absolute vacuum with energy $E_o = E_{vac} = 0$. This leads to

$$V_{2g}(x) = \frac{\alpha^2 \hbar^2 \tilde{E}^2}{2\tilde{m}^3} \nabla^2 w(x) \frac{1}{w(x)} \ . \tag{11}$$

In spherical coordinates this potential is given by

$$V_{2g}(r) = \frac{\alpha^2 \hbar^2 \tilde{E}^2}{2\tilde{m}^3} \ \Big(\frac{d^2 w(r)}{dr^2} + \frac{2}{r}\frac{dw(r)}{dr}\Big) \frac{1}{w(r)} \ . \tag{12}$$

Interestingly, the dynamically generated potential $V_{2g}(r)$ allows a production of $q\bar{q}$-pairs out of the vacuum of fluctuating boson fields, if two boson fields overlap. As shown below, this new type of potential can be identified with the long sought "confinement" potential, required in hadron potential models [12]. The question of a vector or scalar structure of the confinement potential can be studied, see ref. [11], by looking at the splitting of p-wave states in charmonium and bottonium. From the existing data a mixing of vector and scalar structure is found, consistent with the structure of the Lagrangian (6).

For $\mathcal{L}_{3g}$ we obtain a matrix element of the form

$$\mathcal{M}_{3g} = \frac{-\alpha^3}{\tilde{m}} \bar{\psi}(\tilde{p}')\gamma_\mu \; w(q) \frac{g_{\mu\rho} f(p_i)}{{p_i}^2} w(q) \; \gamma_\rho \psi(\tilde{p}) \tag{13}$$

with the condition $\tilde{p}' - \tilde{p} = q + p_i = P$. $f(p_i)$ is the probability to combine P and q to $-p_i$ and has three important consequences: First, it damps out divergencies when $p_i \to 0$: in this case $q \to P$, which is not very likely compared to all other momentum combinations, and we expect $f(p_i) \to 0$. Second, it damps the matrix element for $P \to 0$. This situation requires $q \to -p_i$, which is also suppressed and leads to a finite value of $\mathcal{M}_{3g}$. Finally, it damps out divergencies for $p_i \to \infty$. This can be seen in the Fourier transform of the corresponding potential in r-space discussed below. These arguments indicate that the Lagrangian (7) does not lead to divergencies in the boson-exchange matrix element for $p_i \to 0$ and ∞ and therefore yields solutions in full space.

Contracting the γ-matrices as above and reducing eq. (13) to three dimensions, it can be written by

$$\mathcal{M}_{3g} = \bar{\psi}(\tilde{p}') \; V_{3g}(P) \; \psi(\tilde{p}) \; , \tag{14}$$

where $V_{3g}(P)$ is a Coulomb like potential of the form

$$V_{3g}(P) = \frac{-\alpha^3}{\tilde{m}} \; w(q)^2 \; \frac{f(p_i)}{(p_i)^2} \; . \tag{15}$$

As compared to the Coulomb potential from standard gauge theory, eq. (15) has α^3 coupling, but no divergencies for $p_i \to 0$ and ∞.

Fourier transformation of $V_{3g}(P)$ (15) leads to a folding potential

$$V^v_{3g}(r) = \frac{\hbar}{\tilde{m}} \int dr' \rho(r') \; V_g(r - r') \; , \tag{16}$$

where $\rho(r)$ is a two-boson (quasi) density[3] of a vector state (Fourier transform of the square of the two-boson momentum distribution $w(q)$: $\rho(r) = 2\pi/(\tilde{m}^2\hbar^2) \int pdp \; j_o(pr) \; w(p)^2$) and

---

[3] $\rho(r)$ has dimension fm^{-2} due to the dimension GeV^2 of the two-boson field.

$V_g(r)$ an effective boson-exchange interaction given in r-space by $V_g(r) = -\alpha^3\hbar\frac{f(r)}{r}$. For a $q\bar{q}$ system in a scalar state ($J^\pi = 0^+$), angular momentum L=1 is needed. Therefore, for this case a p-wave density is required in eq. (16), which is related to $\rho(r)$ by

$$\rho^p(\vec{r}) = \rho^p(r)\ Y_{1,m}(\theta,\Phi) = (1+\beta R\ d/dr)\rho(r)\ Y_{1,m}(\theta,\Phi)\ . \tag{17}$$

βR is determined from the condition $< r_{\rho^p} > = \int d\tau\ r\rho^p(r) = 0$ (elimination of spurious motion). This yields a boson-exchange potential given by

$$V^s_{3g}(r) = \frac{\hbar}{\tilde{m}}\int d\vec{r}\,'\rho^p(\vec{r}\,')\ Y_{1,m}(\theta',\Phi')\ V_g(\vec{r}-\vec{r}') = 4\pi\frac{\hbar}{\tilde{m}}\int dr'\rho^p(r')\ V_g(r-r')\ . \tag{18}$$

Further, a geometrical boundary condition, probably related to the field symmetry (1), is assumed, which restricts boson-exchange to the overlap density $\rho(r)$ of two boson fields

$$V^s_{3g}(r) = c_{pot}\ \rho(r)\ , \tag{19}$$

where c_{pot} is a proportionality factor (with dimension $GeV\,fm^3$). A consequence if this is that $< r^2_{V_{3g}} >=< r^2_\rho >$. This boundary condition allows to determine the form of $\rho(r)$, further, it requires that $V_g(r)$ is finite for all values of r (this was already concluded from the combination of the different boson momenta in eq. (13)). To achieve this, a form of the boson-exchange interaction was used

$$V_g(r) = f_{as}(r)(-\alpha^3\hbar/r)\ e^{-cr} \tag{20}$$

with $f_{as}(r) = (e^{(ar)^\sigma}-1)/(e^{(ar)^\sigma}+1)$. As $f_{as}(r) \to 0$ for $r \to 0$ and $e^{-cr} \to 0$ for $r \to \infty$ there are no divergences in $V_g(r)$. A transformation of $f_{as}(r)$ to momentum (Q) space leads to $f_{as}(Q) \to 0$ for $Q \to \infty$. Interestingly, a similar behaviour of the interaction is needed to understand the high momentum behaviour of the strong interaction. In quantum chromodynamics (QCD) this is realized by a decreasing coupling $\alpha(Q)$ related to the property of asymptotic freedom [13].

With the above constraint (19) the self-consistent two-boson density is obtained of the form

$$\rho(r) = \rho_o\ [exp\{-(r/b)^\kappa\}]^2\ \ with\ \ \kappa \sim 1.3 - 1.5\ . \tag{21}$$

This density should be related to the two-boson wavefunction $w(r)$ in the Hamiltonian, see eq. (10), by $\rho(r) = w^2(r)$. Inserting this density in eq. (12) yields the explicite form of the confinement potential

$$V_{2g}(r) = \frac{\alpha^2\hbar^2\tilde{E}^2}{2\tilde{m}^3} \left[\frac{\kappa}{b^2}(\frac{r}{b})^{\kappa-2} \left[\kappa(\frac{r}{b})^{\kappa} - (\kappa+1)\right]\right] . \qquad (22)$$

The mass of the system may be defined in our model by the energy to balance binding (this is also the way hadron masses are observed). This yields

$$M_n = -E_{3g} + E_{2g}^{\ n} \ , \qquad (23)$$

where E_{3g} and $E_{2g}^{\ n}$ are the binding energies in $V_{3g}^s(r)$ or $V_{3g}^v(r)$ and $V_{2g}(r)$, respectively, calculated by using a mass parameter $\tilde{m} = 1/2\ \tilde{M}$. In principle, $\tilde{M}$ should be the mass weighted over different states of the system. However, since this weighting is not known, $\tilde{m} = 1/2\ M$ is used, where M is the ground state mass of the system. This allows to use the additional constraint $M = M_1$. The mass contributions due to excited states is included in $\tilde{E}$, which is used as fit parameter.

Concerning the binding energies, E_{3g} is negative. Since the $q\bar{q}$ pairs are massless, the averaged energy over different states $\tilde{E}_{3g}$ can be related to the mean square momentum $< Q^2_\rho >$ of $\rho(r)$ by the energy-momentum relation $E_{vac} = 0 = \sqrt{< Q^2_\rho >} + \tilde{E}_{3g}$. This gives

$$\tilde{E}_{3g} = -\sqrt{< Q^2_\rho >} \ , \qquad (24)$$

which is used as another constraint in the calculations. The binding in $V_{3g}(r)$ does not correspond to real mass generation. However, the binding energy E_{2g} is positive. This allows creation of stable particles out of the absolute vacuum of fluctuating boson fields, if two rapidly fluctuating boson fields overlap and trigger a quantum fluctuation with energy E_{2g}.

An effective coupling constant α^3 of 1/137 is used for atomic systems, whereas for the hadronic case α is determined by matching the mass to the lowest binding energy in eq. (23).

As a test case, an application of the above formalism is discussed for the lowest bound states in the hydrogen atom $e^- - p$ and positronium $e^+ - e^-$, which have been studied in very different approaches, the Bohr model, non-relativistic Schrödinger equation, relativistic Dirac equation and different QED methods. For atomic systems the q-q bound

Table 1: Results for $e^- - p$ and $e^+ - e^-$ bound states. Binding energies, $\tilde{m}$ and average Q in eV, root mean square radius and b in pm, c and a in pm^{-1}.

	System	E_{3g}(scalar)	E_{3g}(vector)	$E_{exp}(1s)$	$E_{exp}(2p)$	$< r^2 >^{1/2}$	R_{Bohr}
1.	$e^- - p$	-13.6	-3.4	-13.6	-3.4	57	53
2.	$e^+ - e^-$	-6.8	-1.7	-6.8	-1.7	114	106

	κ	b	c	a	σ	α^3	$\tilde{m}$	$\tilde{E}$	$< r_\rho^2 >^{1/2}$
1.$^{*)}$	1.3	$0.14\ 10^5$	$1.2\ 10^{-4}$	$1.2\ 10^{-3}$	0.86	1/137	6.9	0.16	16500
final		$0.47\ 10^2$	$3.8\ 10^{-2}$	$1.6\ 10^{-1}$	0.86				57
2.$^{*)}$	1.3	$0.24\ 10^5$	$0.7\ 10^{-4}$	$0.7\ 10^{-3}$	0.86	1/137	3.4	0.08	28700
final		$0.94\ 10^2$	$2.0\ 10^{-2}$	$0.8\ 10^{-1}$	0.86				114

$^{*)}$ parameters without recoil mass correction.

state energies are much smaller than the masses of the participating particles. Therefore, the potentials $V_{3g}(r)$, given by eqs. (16) and (18) and $V_{2g}(r)$ (22) have to be modified by the reduced mass $\mu = m_1 m_2/(m_1 + m_2)$ of the system (with $m_1 = m_e$ and $m_2 = m_p$ or m_e for the two cases considered). In the present generalised form of QED no further corrections have to be applied.

First, interaction potentials (16) and (18) for scalar (1s) and vector (2p) q-q states were calculated (without recoil corrections due to the masses m_1 and m_2). The slope parameter b was adjusted to give the known binding energies of the 2p-state. The resulting interaction parameters c, a, σ and α^3 are given in table 1. The differences in the parameters between the $e^- - p$ and $e^+ - e^-$ system are entirely due to the fact that the reduced mass μ differs by a factor of two.

Results for the $e^- - p$ bound state system are given in fig. 1. The low-radius cut-off function $f_{as}(r)$ and the relative boson-exchange interaction $V_g(r)$ are shown in the upper part by the dashed and solid line, respectively. A comparison of the boson-exchange interaction with the Coulomb interaction, given the dot-dashed line, shows only deviations at small and large radii, indicating that the resulting spectrum does not deviate much from that generated by the Coulomb potential. The boson-exchange potential is finite for $r \to 0$ and ∞, therefore, solutions exist in the infrared and ultraviolet momentum region and do not require non-relativistic or perturbative approximations.

In the middle part of fig. 1 a comparison of the two-boson density (dot-dashed line) with the boson-exchange potential (solid line) is made. One can see that by the choice of parameters in table 1 both quantities are in close agreement, as required from relation (19). The 'confinement' potential $V_{2g}(r)$ is shown in the lower part, which has a radial structure very different from the boson-exchange potential $V_{3g}(r)$. It is very flat and can be well approximated at larger radii by a linear form. Due to its flatness and the fact that it has negative and positive values in different regions of r, the corresponding bound state energy E_{2g} is only 0.4 10^{-3} eV. Therefore, its contribution to the binding energy of the system is completely negligible.

The resulting binding energies for the 1s and 2p states are given in table 1. They are in good agreement with the known binding energies [14]. Also the binding energies for the 2s state of -3.4 eV and 3s and 3p states of -1.51 eV for hydrogen and the corresponding binding energies of -1.7 and -0.755 eV for positronium are consistent with experiment. The factor two between hydrogen atom and positronuim is due to the different reduced masses of the two systems. Consequently, in the first case the root mean square radius is smaller by a factor 2 (with some numerical uncertainty).

In a second step, the recoil due to the $p + e^-$ and $e^+ + e^-$ masses has been taken into account. This has been done by modifying the radial densities, using a mass parameter $\tilde{m} = 1/2\ \tilde{M} = 1/2(\mu - E_{3g} + E_{2g})$, with the reduced mass μ defined above. The resulting parameters are also given in table 1 with the label 'final'. For the $e^- + p$ system the corresponding two-boson density and boson-exchange potential are shown in the upper part of fig. 2 by dot-dashed and solid line, respectively, which are strongly compressed in radius with respect to the same quantities in fig. 1. Again one can see close agreement between $\rho(r)$ and $V^s_{3g}(r)/c_{pot}$, as required from relation (19). In the middle part of fig. 2 the Fourier tansformed quantities are given. In transforming the two-boson density and boson-exchange potential (18) to Q-space agreement is obtained only, if the process $gg \rightarrow q\bar{q}$ is elastic and consequently the created $q\bar{q}$-pair has no mass. Thus, our formalism requires an interaction between massless fermions (quantons). The recoil compressed 'confinement' potential is given in the lower part of fig. 2. Here the contribution to the mass is enhanced but still two orders of magnitude smaller than the contribution from $V_{3g}(r)$.

Of interest is also the radial extent of the bound state systems in question. The root mean square radius for both systems is very close to the Bohr radius (see table 1). However, we see in the upper part of fig. 2 that the density falls off quite smoothly and is already quite reduced at the rms radius. Therefore, a much smaller radius at full width half maximum R_{fwhm} of 20 pm and 40 pm is extracted for system 1 and 2, respectively. Interestingly, the average radius $\tilde{R} = \int \rho(r) r^2 dr / \int \rho(r) r dr$ of 36 pm for the hydrogen atom is in good agreement with the covalent radius of 31±5 pm [15].

A second evaluation of the above formalism is discussed for the lowest scalar mesonic system $\sigma(600)$, which has only recently been settled experimentally [16]. For hadrons usually QCD is applied, however, the fact that hadrons are colour neutral and the correct confinement potential [12] is obtained in our formalism (as seen below) suggests that a treatment in the quanton model is justified. A detailed comparison with QCD will be discussed in a forthcoming paper.

Results on the effective interaction, the two-boson density and the boson-exchange potential are given in fig. 3 and table 2. Similar to the first case, the slope parameter b has been adjusted to give the mass of the lowest scalar state. In the upper part of fig. 3 the low radius cut-off function $f_{as}(r)$ and the effective boson-exchange interaction is given by the dashed and solid lines. We see a quite similar behaviour as for the atomic system in fig. 1. In the two lower parts a comparison of the two-boson density (dot-dashed lines) with the boson-exchange potential (solid lines) is made. By the choice of parameters in table 2 both quantities are again in close agreement, as required from relation (19). Also in the Fourier transformation of two-boson density and boson-exchange potential to Q-space agreement can be obtained only, if the elementary fermions are massless.

However, very different from the atomic systems the 'confinement' potential $V_{2g}(r)$ is much stronger, although the relative shape is quite similar. This is shown in fig. 4. It has the same form as the known confinement potential $V_{conf} = -\alpha/r + l \cdot r$ deduced from potential models [12] and the lattice data of Bali et al. [17] (solid points). Further, this potential is similar for other solutions of smaller or larger radii and can therefore be identified with the 'universal' confinement potential in hadrons. It is important to note, that the potential (22) reproduces the 1/r + linear form without any assumption on its distance behavior; this is entirely a consequence of the deduced radial form (21) of the two-boson density.

Table 2: Results for $\sigma(600)$ and higher excited scalar mesons in comparison with known 0^{++} mesons [1]. Masses and average Q in GeV, mean radius square in fm^2. The parameters given in the lower part (b in units of fm, c and a in units of fm^{-1}) are constrained by the relations (19), (23) and (24). For the lowest state the obtained energies E_{2g} and E_{3g} are 0.4 GeV and -0.15 GeV, respectively.

Meson)	M_1	M_2	M_3	M_1^{exp}	M_2^{exp}	$< r_\rho^2 >$	$< Q_\rho^2 >^{1/2}$
$\sigma(600)$	0.55	1.28	1.88	0.55±0.2	1.35±0.2	0.65	0.59

κ	b	c	a	σ	α	$\tilde{m}$	$\tilde{E}$
1.5	0.77	2.4	6.35	0.86	0.26	0.14	0.90

Using all constraints discussed above lead to a complete determination of all parameters of the model (with some uncertainties) for a given slope parameter b. The last open parameter b may be determined by the structure of the vacuum. For mesonic systems this is discussed in ref. [18] and leads to a description, in which all parameters of the model are fixed.

The width of the state is given by the derivative of the momentum distribution, but threshold effects have to be included as well. The calculated mass distribution (without threshold effects) is given by the dot-dashed line in fig. 5. Using the threshold function from ref. [19] with parameters M_o=0.55 GeV and Γ=0.55 GeV, a final mass distribution is obtained in good agreement with the shape of the $\sigma(600)$ meson extracted experimentally [16].

Finally, the resulting value of $\tilde{E}$ of about 0.9 GeV is in reasonable agreement with the mean energy of scalar excitation including excited states, see table 2 (which shifts the energy from 0.55 GeV by several tenths of GeV). This can be taken as a further consistency check of our model.

In summary, the inclusion of scalar two-boson coupling in QED leads to new properties of boson-exchange matrix elements, which allow a satisfactory description of basic properties of atomic and hadronic systems. For these very different bound states a large similarity in the relative shape of boson-exchange potentials is observed, but large differences in the dynamically generated confinement potential. This is consistent with the observation that hadrons are confined but light atoms have only small binding energies of a few eV. The fact that small and large systems are described equally well, not dependent on fine-tuning, shows that the model is robust, important for a description of different fundamental forces. It appears that the present form of the Lagrangian (giving rise to static properties of bound states as well as dynamical features, which are missing in the SM and other extensions of it) is consistent with the general laws of nature, which require imperatively static as well as dynamic components for evolution.

We are very grateful to P. Decowski, M. Dillig (deceased), A. Kupsc, and P. Raczka among many other colleagues for fruitful discussions, valuable comments and the help in formal derivations. Special thanks to P. Zupranski for numerous conversations and B. Loiseau for his help with the formulation of the Lagrangian.

References

[1] Review of particle properties, K. Nakamura et al., J. Phys. G 37, 075021 (2010); http://pdg.lbl.gov/ and refs. therein

[2] G. Aad et al. Phys. Rev. Lett. 106, 131802 (2011); see also
D. Toback, APS-J. (Part. and Fields) Phys. 4, 27 (2011) and refs. therein

[3] H.P. Morsch and P. Zupranski, Phys. Rev. C 71, 065203 (2005) and refs. therein

[4] H.P. Morsch, Z. Phys. A 350, 61 (1994)

[5] R. Machleidt, Adv. Nucl. Phys. 19 (1989) 189; and refs. therein

[6] H.P. Morsch, W. Spang and P. Decowski, Phys. Rev. C 67, 064001 (2003)

[7] A. Donnachie and P.V. Landshoff, Nucl. Phys. B 231 (1984) 189 and B 311 (1989) 509; P.V. Landshoff and O. Nachtmann, Z. Phys. C 35, 405 (1987); F. Halzen, G. Krein and A.A. Natale, Phys. Rev. D 47, 295 (1993)

[8] J.Z. Simon, Phys. Rev. D 41, 3720 (1990); A. Foussats, E. Manavella, C. Repetto, O.P. Zandron, and O.S. Zandron, Int. J. theor. Phys. 34, 1 (1995); V.V. Nesterenko, J. Phys. A: Math. Gen. 22, 1673 (1989); and refs. therein

[9] In the matrix elements (8) and (13) the indices of the γ-matrices are arbitrary; so, there is no difference, if the combinations $\gamma_\mu\gamma_\rho$ or $\gamma_\rho\gamma_\mu$ are chosen.

[10] The two-boson field $w(q)$ yields contributions only, if the time-components of both overlapping boson fields are equal. This allows a reduction of the matrix elements to three space dimensions.

[11] H.J. Schnitzler, Phys. Rev. Lett. 35, 1540 (1975)

[12] R. Barbieri, R. Kögerler, Z. Kunszt, and R. Gatto, Nucl. Phys. B 105, 125 (1976); E. Eichten, K.Gottfried, T. Kinoshita, K.D. Lane, and T.M. Yan, Phys. Rev. D 17, 3090 (1978); S. Godfrey and N. Isgur, Phys. Rev. D 32, 189 (1985); D. Ebert, R.N. Faustov, and V.O. Galkin, Phys. Rev. D 67, 014027 (2003); and refs. therein

[13] D.J. Gross and F. Wilczek, Phys. Rev. Lett. 30, 1343 (1973); H.D. Politzer, Phys. Rev. Lett. 30, 1346 (1973)

[14] S and p state energies are calculated in the potentials $V^s_{3g}(r)$ and $V^v_{3g}(r)$ using different densities, see eq. (17). The fact that similar binding energies are deduced as for a simple Coulomb potential supports the correctness of the used densities.

[15] B. Cordera et al., Covalent radii revisited, Daton Trans. 21, 2832 (2008)

[16] M. Ablikim, et al., hep-ex/0406038 (2004); see also D.V. Bugg, hep-ex/0510014 (2005) and refs. therein

[17] G.S. Bali, K. Schilling, and A. Wachter, Phys. Rev. D 56, 2566 (1997);
G.S. Bali, B. Bolder, N. Eicker, T. Lippert, B. Orth, K. Schilling, and T. Struckmann, Phys. Rev. D 62, 054503 (2000)

[18] H.P. Morsch, arXiv 1104.2575 [hep-ph], with Lagrangian to be revised, see eq. (3). However, all calculations are based on the potentials $V_{2g}(r)$ and $V_{3g}(r)$ and lead therefore to unchanged results.

[19] H.P. Morsch and P. Zupranski, Phys. Rev. C 61, 024002 (1999)



Figure 1: Self-consistent solution for $e^- - p$ bound state system (without p and e^- masses taken into account). Upper part: Low radius cut-off function $f_{as}(r)$ (dashed line) and relative shape of the interaction (20) given by solid line in comparison with a pure Coulomb interaction (dot-dashed line). Middle part: Two-boson density and boson-exchange potential $|V^s_{3g}(r)/c_{pot}|$ given by the overlapping dot-dashed and solid lines, respectively. Lower part: 'Confinement' potential $V_{2g}(r)$.



Figure 2: Self-consistent solution for $e^- - p$ bound state system (including p and e^- masses). Upper two parts: Two-boson density and boson-exchange potential $|V^s_{3g}/c_{pot}|$ in r- and Q-space given by the overlapping dot-dashed and solid lines, respectively. Lower part: 'Confinement' potential $V_{2g}(r)$.



Figure 3: Self-consistent solution for a scalar state with mass=0.55 GeV and $< r_\rho^2 >$=0.65 fm^2. Upper part: Low radius cut-off function $f_{as}(r)$ and relative shape of the interaction (20) given by dashed and solid line, respectively. Lower two parts: Two-boson density and boson-exchange potential $|V_{3g}^s/c_{pot}|$ given by the overlapping dot-dashed and solid lines, respectively, in r- and Q-space.



Figure 4: Deduced 'confinement' potential $V_{2g}(r)$ (solid line, multiplied by a factor of 3.7) in comparison with the data from lattice gauge calculations [17].



Figure 5: Mass distribution (dot-dashed) and calculated spectral shape (solid line) of the scalar state with a mass of 0.55 GeV in comparison with the experimental results on the $\sigma(600)$ meson from Bugg et al. [16].